

Buffer and Media Preparation – What Is a Buffer?

Your Objectives:

At the end of the lesson you should be will able to explain how to make a buffer solution.

A **buffer** (more precisely, pH buffer or hydrogen ion buffer) is an solution consisting of a mixture of a weak and its conjugate base, or vice versa). Its pH changes insignificantly when a small amount of strong acid or is added to it. Buffer solutions are used as a means of keeping pH at a nearly constant value in a wide variety of applications. In nature, there are many systems that use buffering for pH . For example, the bicarbonate buffering system is used to the pH of blood.

pH

In , pH (denoting 'potential of or 'power of hydrogen') is a scale used to specify the or of an aqueous solution. Lower pH correspond to solutions which are more acidic in nature, while higher values correspond to solutions which are more basic or . At room temperature (25 C or 77 F), pure water is neutral (neither acidic nor basic) and has a pH of 7.

Substance	pH range	Type
Battery acid	< 1	Acid
Gastric acid	1.0 – 1.5	
Vinegar	2.5	
Orange juice	3.3 – 4.2	
Black coffee	5 – 5.03	
Milk	6.5 – 6.8	Neutral
Pure water	7	
Sea water	7.5 – 8.4	Base
Ammonia	11.0 – 11.5	
Bleach	12.5	
Lye	13.0 – 13.6	

The pH is logarithmic and inversely indicates the concentration of ions in the (a lower pH indicates a higher concentration of hydrogen ions). More precisely, pH is the negative of the base-10 logarithm of the activity of the hydrogen ion.


At 25°C, solutions with a pH less than 7 are , and solutions with a pH

greater than 7 are . The value of the pH

depends on the temperature being lower than 7 if the temperature increases. The pH value can be less than 0 for very strong acids, or greater than 14 for very strong bases. The pH scale is traceable to a set of standard solutions whose pH is established by international

.


Buffer solutions achieve their to pH change because of the presence of an between the weak acid HA and its conjugate base A⁻:



When some strong acid is added to an equilibrium mixture of the weak acid and its conjugate base, hydrogen ions (H⁺) are added, and the equilibrium is shifted to the left, in accordance with Le Châtelier's . Because of this, the hydrogen ion


increases by less than the amount expected for the quantity of strong acid added. Similarly, if strong alkali is added to the mixture, the hydrogen ion concentration decreases by less than the amount expected for the quantity of alkali added. The

is illustrated by the simulated titration of a weak acid with pK_a = 4.7. The relative concentration of undissociated acid is shown in blue, and of its base in red.




The changes relatively slowly in the

region, $\text{pH} = \text{pK}_a \pm 1$, centered at $\text{pH} = 4.7$, where $[\text{HA}] = [\text{A}^-]$. The hydrogen ion concentration decreases by less than the amount expected because most of the added hydroxide ion is consumed in the .



and only a little is consumed in the reaction (which is the reaction that results in an increase in pH)



Once the acid is more than 95% deprotonated, the pH rapidly since most of the added alkali is consumed in the neutralization reaction.

Helpful link: <https://www.khanacademy.org/science/ap-chemistry/buffers-titrations-solubility-equilibria-ap/buffer-solutions-tutorial-ap/v/buffer-system>

The pH of a solution containing a buffering can only vary within a narrow , regardless of what else may be present in the solution. In

systems this is an essential condition for

to function correctly. In human blood, for instance, we find a mixture of carbonic acid (H_2CO_3) and bicarbonate (HCO_3^-) present in the plasma fraction, which constitutes the major mechanism for maintaining the level of blood at between 7.35 and 7.45. Outside this narrow range (7.40 ± 0.05 pH unit), acidosis and alkalosis metabolic conditions rapidly develop, ultimately leading to death if the correct buffering capacity is not rapidly restored.

If the pH value of a rises or falls too much, the effectiveness of an enzyme decreases in a process, known as , which is usually irreversible. The majority of biological that are used in research are kept in a buffer solution, often phosphate buffered saline (PBS) at pH 7.4.

In industry, agents are used in fermentation processes as well as in setting the correct conditions for dyes used in colouring fabrics. They are also used in chemical and of pH meters.

Simple buffering agents

Buffering agent	Useful pH range
Citric acid	2.1–7.4
Acetic acid	3.8–5.8
KH_2PO_4	6.2–8.2
CHES	8.3–10.3
Borate	8.25–10.25

For buffers in acidic , the pH may be to a desired value by adding a acid, such as hydrochloric acid, to the particular buffering agent. For alkaline buffers, a strong base, such as sodium hydroxide, may be added. Alternatively, a buffer can be made from a mixture of an acid and its conjugate base. For example, an acetate buffer can be made from a mixture of acetic acid and sodium acetate. Similarly, an alkaline buffer can be made from a mixture of the base and its conjugate acid.

Aufgabe Lückentext:

Folgende Wörter bitte in den Lückentext einfüllen.

Jedes Wort kommt einmal vor.

Bitte Gross- und Kleinbuchstaben beachten.

aqueous, analysis, agreement, acidity, acid, acidic, adjusted, agent, alkaline, base, basicity, basic, biological, buffer, buffering, calibration, chemical, chemistry, concentration, conjugate, denaturation, equilibrium, effect, enzymes, hydrogen', hydrogen, mixture, neutral, neutralization, principle, pH, pH, range, regulation, regions, regulate, reaction, resistance, rises, samples, solution, solution, solution, scale, strong, values,